

*Pacific
Journal of
Mathematics*

**UNCOUNTABLY MANY INEQUIVALENT LIPSCHITZ
HOMOGENEOUS CANTOR SETS IN \mathbb{R}^3**

DENNIS GARITY, DUŠAN REPOVŠ AND MATJAŽ ŽELJKO

UNCOUNTABLY MANY INEQUIVALENT LIPSCHITZ HOMOGENEOUS CANTOR SETS IN \mathbb{R}^3

DENNIS GARITY, DUŠAN REPOVŠ AND MATJAŽ ŽELJKO

General techniques are developed for constructing Lipschitz homogeneous wild Cantor sets in \mathbb{R}^3 . These techniques, along with Kauffman's version of the Jones polynomial and previous results on Antoine Cantor sets, are used to construct uncountably many topologically inequivalent such wild Cantor sets in \mathbb{R}^3 . This use of three-dimensional finite link invariants to detect distinctness among wild Cantor sets is unexpected. These Cantor sets have the same Antoine graphs and are Lipschitz homogeneous. As a corollary, there are uncountably many topologically inequivalent Cantor sets with the same Antoine graph.

1. Introduction

Malešič and Repovš [1999] have constructed a specific example of a wild Cantor set in \mathbb{R}^3 that is Lipschitz homogeneously embedded. This answered negatively a question in [Repovš et al. 1996] as to whether Lipschitz homogeneity of a Cantor set implied tameness. In this paper, we introduce more general techniques for detecting the Lipschitz homogeneity of Cantor sets in \mathbb{R}^n . These techniques allow us to construct uncountably many topologically distinct Lipschitz homogeneous wild Cantor sets in \mathbb{R}^3 . These Cantor sets are all simple Antoine Cantor sets with the same Antoine graph as defined in [Wright 1986]. The fact that the constructed Cantor sets are all topologically distinct is a consequence of a result of Sher [1968] and a computation of Kauffman's version [1988] of the Jones polynomial for the center lines of certain tori used in the construction. It is hoped that the techniques in this paper may also prove to be applicable to showing that certain Blankinship type Cantor sets [Blankinship 1951; Eaton 1973] in \mathbb{R}^n for $n \geq 4$ can be constructed so as to be Lipschitz homogeneously embedded.

MSC2000: primary 54E45, 54F65; secondary 57M30, 57N10.

Keywords: wild Cantor set, Lipschitz homogeneity, similitude, coefficient of similarity, defining sequence, link invariant.

Garity was supported in part by NSF grants DMS 0139678 and DMS 0104325. Repovš and Matjaž were supported in part by MESS grant 0101-509. All authors were supported in part by MESS grant SLO-US 2002/01.

2. Notation and Background

Lipschitz maps and similitudes. A map $S: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be a *Lipschitz map* if there exists a constant λ such that

$$|S(x) - S(y)| \leq \lambda|x - y| \text{ for every } x, y \in \mathbb{R}^n$$

and the smallest such λ is called the *Lipschitz constant of S* . In the special case when

$$|S(x) - S(y)| = \lambda|x - y| \text{ for every } x, y \in \mathbb{R}^n$$

the map S is called a *similarity* and the number λ is called the *coefficient of similitude*. Finally, when $\lambda = 1$ the map S is called an *isometry*.

A Cantor set C in \mathbb{R}^3 is *Lipschitz homogeneously embedded* if for each pair of points x and y in C there is a Lipschitz homeomorphism $h: \mathbb{R}^n \rightarrow \mathbb{R}^n$ with h^{-1} also Lipschitz such that $h(C) = C$ and $h(x) = y$.

Coordinates of points in Cantor sets. In the applications of this section, the compact set X mentioned below will in general be a solid torus.

Let $G_i, 1 \leq i \leq M$, be finite index sets and let $\mathbf{S}_i = \{S_g: \mathbb{R}^n \rightarrow \mathbb{R}^n : g \in G_i\}$ be a set of similarities having the same coefficient λ_i of similitude. Let $\mathbf{S} = \cup \mathbf{S}_i$. Additionally, suppose that there exists a compact set $X \subset \mathbb{R}^n$ such that

- (1) $S_g(X) \subset \overset{\circ}{X}$ for each $g \in G_i$ and
- (2) the sets $S_g(X)$ are pairwise disjoint, $g \in G_i$.

Let $\mathbf{T} = (n_1, n_2, \dots)$ be a fixed sequence where each n_i is in $\{1, \dots, M\}$. Let $G^k = G_{n_1} \times G_{n_2} \times \dots \times G_{n_k}, G_k^\infty = \prod_{i=k}^\infty G_{n_i}$ and $G^\infty = \prod_{i=1}^\infty G_{n_i}$. For each multiindex $\gamma = (g_1, g_2, \dots, g_k) \in G^k$, write

$$S_\gamma = S_{g_1} \circ S_{g_2} \circ \dots \circ S_{g_k} \quad \text{and} \quad X_\gamma = S_\gamma(X).$$

In particular, $X_g = S_g(X)$ for $g \in G$.

The number of components of a multiindex γ is called the *depth of γ* . So depth $\gamma = k$ if $\gamma \in G^k$.

$$\text{Let } X_k = \bigcup_{\text{depth } \gamma=k} X_\gamma.$$

It is well-known [Hutchinson 1981] that the intersection of the sequence of sets $X \supset X_1 \supset X_2 \supset \dots$ is a Cantor set. Denote this set by $|(S, T)|$. This Cantor set is self-similar if \mathbf{T} is repeating.

For an infinite multiindex $\gamma = (g_1, g_2, g_3, \dots) \in G^\infty$ define

$$\gamma^k = (g_1, g_2, \dots, g_k) \quad \text{and} \quad X_\gamma = \bigcap_{k=1}^\infty X_{\gamma^k}.$$

Obviously, each X_γ is a singleton, consisting of a point from the Cantor set $|(\mathbf{S}, \mathbf{T})|$ and for each point from $|(\mathbf{S}, \mathbf{T})|$ there exists exactly one such multiindex γ . The components of γ are called the *coordinates* of the corresponding point from the Cantor set $|(\mathbf{S}, \mathbf{T})|$.

Finally define a juxtaposition of multiindices as follows. If $\delta = (d_1, d_2, \dots, d_k)$ is a finite multiindex and $\gamma = (g_1, g_2, \dots)$ is finite or infinite then let

$$\delta\gamma = (d_1, d_2, \dots, d_k, g_1, g_2, \dots).$$

In the special case when depth $\gamma = 1$, we have $\gamma = g_1$ and $\delta g_1 = (d_1, d_2, \dots, d_k, g_1)$.

Antoine Cantor sets. We give a brief summary of results from [Sher 1968] and [Wright 1986]. Sher's results are needed in our proof of our main theorem, while Wright's are used in our observation about Antoine graphs associated with the Cantor sets we construct.

An *Antoine Cantor set* C in \mathbb{R}^3 is a Cantor set meeting the following conditions.

- (1) C has a defining sequence M_1, M_2, \dots , each M_i consisting of the union of a finite number of pairwise disjoint standard unknotted solid tori in \mathbb{R}^3 and M_1 consisting of a single solid torus.
- (2) The tori in M_i , for $i \geq 2$, can be listed in a sequence $T_{i,1}, T_{i,2}, \dots, T_{i,n(i)}$ so that T_j and T_k are of simple linking type if $j - k = \pm 1 \pmod n$ and do not link if $j - k \neq \pm 1 \pmod n$.
- (3) The linked chain of tori $T_{i,1}, T_{i,2}, \dots, T_{i,n(i)}$ have winding number greater than 0 in the torus at the previous stage that contains them.


If, in condition (3), the winding number is required to be 1, and if each $n(i) \geq 4$, we call the resulting Cantor set a *simple Antoine Cantor set*. Most Antoine Cantor sets in the literature, including the original one [Antoine 1920] are simple.

Sher [1968] showed that if two Antoine Cantor sets C_1 and C_2 with defining sequences M_1, M_2, \dots and N_1, N_2, \dots are equivalently embedded in \mathbb{R}^3 , then there is a homeomorphism h of \mathbb{R}^3 to itself such that for each i , h takes the tori in M_i homeomorphically onto the tori in N_i . As a consequence, if it can be shown that for some i , no such homeomorphism exists, the two Cantor sets are inequivalently embedded. This is the result we will need to construct the uncountably many inequivalently embedded Cantor sets.

Wright [1986] associates an Antoine graph $\Gamma(C)$ with a Antoine Cantor C with defining sequence M_1, M_2, \dots . The graph Γ is a countable union of nested subgraphs $\Gamma_0 \subset \Gamma_1 \subset \Gamma_2 \subset \dots$. The subgraph Γ_0 is a single vertex. For each vertex v of $\Gamma_i - \Gamma_{i-1}$, there is a polygonal simple closed curve with at least 4 vertices $P(v)$ contained in $\Gamma_{i+1} - \Gamma_i$ so that if v and w are distinct vertices of $\Gamma_i - \Gamma_{i-1}$, then $P(v)$ and $P(w)$ are disjoint. Γ_{i+1} consists of Γ_i , the union of the $P(v)$ for v

in Γ_i , and the union of edges running from v to the vertices of $P(v)$. The vertices of $\Gamma_i - \Gamma_{i-1}$ correspond to the components of M_i .

If v is a vertex of $\Gamma_i - \Gamma_{i-1}$ corresponding to a component C of M_i , then $P(v)$ contains precisely the vertices corresponding to the components of M_{i+1} contained in C , and two such vertices are joined by an edge if and only if the corresponding components of M_{i+1} are linked. Here is a diagram of an Antoine graph similar to that in [Wright 1986, p. 252]:



Wright shows that if C_1 and C_2 are simple Antoine Cantor sets with different Antoine graphs $\Gamma(C_1)$ and $\Gamma(C_2)$, the Cantor sets are inequivalently embedded.


In our construction, all of the Cantor sets constructed have the same Antoine graph, but are inequivalently embedded.

3. Constructing Lipschitz homogeneous Cantor sets

Let G_i , $1 \leq i \leq M$, $\mathbf{S}_i = \{S_g: \mathbb{R}^n \rightarrow \mathbb{R}^n; g \in G_i\}$, X , X_g and $\mathbf{T} = (n_1, n_2, \dots)$ be as above. The setting to keep in mind when reading Theorem 1 below is that of a simple Antoine Cantor set defined by tori where each stage m torus has $|G_{n_m}|$ stage $m + 1$ tori in its interior. For Theorem 1, also assume that each G_i is a finite cyclic group, with the group operation written additively.

Theorem 1. *For each i , $1 \leq i \leq M$, suppose that $f_i: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Lipschitz homeomorphism and that*

- (i) $f_i|_{\mathbb{R}^n - \hat{X}} = \text{id}$,
- (ii) *for each $g \in G_i$, we have $f_i(X_g) = X_{g+1}$ and the diagram*



commutes.

Then $|(S, \mathbf{T})|$ is Lipschitz homogeneous in \mathbb{R}^n .

Proof. The approach to the proof is similar to that used in [Malešič and Repovš 1999, Lemma 1]. The main modification needed is to take into account the presence of more than one finite index set. Fix an arbitrary pair of points a and b in $|(\mathbf{S}, \mathbf{T})|$. We will construct a homeomorphism

$$h : (\mathbb{R}^n, |(\mathbf{S}, \mathbf{T})|, a) \rightarrow (\mathbb{R}^n, |(\mathbf{S}, \mathbf{T})|, b)$$

and prove that both h and h^{-1} are Lipschitz. Let $\alpha = (a_1, \dots, a_k, \dots)$ and $\beta = (b_1, \dots, b_k, \dots) \in G^\infty$ be the coordinates of a and b . For an arbitrary $\gamma = (g_1, \dots, g_k) \in G^k$ define the homeomorphism

$$f_\gamma = S_\gamma \circ f_{n_{k+1}} \circ S_\gamma^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n.$$

Set

$$\begin{aligned} r_1 &= f_{n_1}^{b_1 - a_1}, r_2 = f_{b_1}^{b_2 - a_2}, r_3 = f_{(b_1, b_2)}^{b_3 - a_3}, \dots, r_{k+1} = f_{\beta^k}^{b_{k+1} - a_{k+1}}, \dots \\ \ell_i &= r_i^{-1}, \\ h_k &= r_k \circ r_{k-1} \circ \dots \circ r_2 \circ r_1. \end{aligned}$$

In addition, for notational convenience, let


$$r_{-1} = r_0 = id|_{\mathbb{R}^n}.$$

It follows by Lemma 2(iv) below that the sequences of homeomorphisms h_1, h_2, \dots and $h_1^{-1}, h_2^{-1}, \dots$ converge pointwise at all points different from the point a and b , respectively. The convergence of the sequences at the point a and at the point b follows from Lemma 2(ii). Denote the limits of the sequences by $h : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\tilde{h} : \mathbb{R}^n \rightarrow \mathbb{R}^n$, respectively. It also follows by Lemma 2 that $h(a) = b$, that $h(|(\mathbf{S}, \mathbf{T})|) = |(\mathbf{S}, \mathbf{T})|$, and that $h \circ \tilde{h} = \tilde{h} \circ h = id_{\mathbb{R}^n}$. It follows from Lemma 3 that h and \tilde{h} are Lipschitz. Thus Theorem 1 is proved. \square

Lemmas needed for Theorem 1.

Lemma 1. *The homeomorphism f_γ is Lipschitz with Lipschitz constant equal to that of $f_{n_{k+1}}$. Moreover:*

- (i) $f_\gamma|_{\mathbb{R}^n - \hat{X}_\gamma} = id$.
- (ii) For arbitrary $g \in G_{n_{k+1}}$, we have $f_\gamma(X_{\gamma g}) = X_{\gamma(g+1)}$, the diagram



commutes, and $f_\gamma|_{X_{\gamma g}}$ is an isometry.

(iii) For arbitrary $(g_{k+1}, g_{k+2}, \dots) \in G_{k+1}^\infty$,

$$f_\gamma(X_{(\gamma, g_{k+1}, g_{k+2}, \dots)}) = f_\gamma(X_{(\gamma, 1+g_{k+1}, g_{k+2}, \dots)}).$$

Proof. This follows in a manner similar to [Malešič and Repovš 1999, Lemma 2]. Part (i) follows directly from condition (i) of Theorem 1. Part (ii) follows from condition (ii) of the same theorem. Finally, (ii) implies (iii). \square

Lemma 2. *The homeomorphisms h_k exhibit the following properties:*

- (i) $h_k^{-1} = \ell_1 \circ \ell_2 \circ \dots \circ \ell_{k-1} \circ \ell_k$.
- (ii) $h_k(X_{\alpha^k}) = X_{\beta^k}$ and $h_k(X_{\alpha^k \gamma}) = X_{\beta^k \gamma}$ for arbitrary multiindex γ .
- (iii) the restriction $h_k|_{X_{\alpha^k a_{k+1}}} : X_{\alpha^k a_{k+1}} \rightarrow X_{\beta^k a_{k+1}}$ is an isometry.
- (iv) $h_k|_{\mathbb{R}^n - \hat{X}_{\alpha^k}} = h_{k+1}|_{\mathbb{R}^n - \hat{X}_{\alpha^k}} = h_{k+2}|_{\mathbb{R}^n - \hat{X}_{\alpha^k}} = \dots$.

Proof. This follows in a manner similar to [Malešič and Repovš 1999, Lemma 3]. Property (i) can be proved directly by examining the construction of h_k . Property (ii) follows from Lemma 1(ii)–(iii). Property (iii) holds since $f_\gamma|_{X_{\gamma g_{k+1}}}$ is an isometry. Property (iv) holds because of Lemma 1(i). \square

Lemma 3. *h_k and h_{k-1} are Lipschitz with equal Lipschitz constants for all values of k .*

Proof. This requires the most modification of [Malešič and Repovš 1999] as multiple similarities with different constants of similarity are involved.

We fix the sequence $\alpha = (a_1, a_2, \dots)$ of coordinates of the point $a \in |\mathbf{S}|$ and introduce the notion of *depth* of a point $x \in \mathbb{R}^n$:

$$\text{dep } x = j \quad \text{if } x \in X_{\alpha^j} - \hat{X}_{\alpha^{j+1}}.$$

Additionally, let

$$\text{dep } x = 0 \quad \text{if } x \in X - \hat{X}_{\alpha_1} \quad \text{and} \quad \text{dep } x = -1 \quad \text{if } x \in \mathbb{R}^n - \hat{X}.$$

In the case $x \in \hat{X}_{\alpha^j}$ for all $j \in \mathbb{N}$ (i.e. $x = a$) let $\text{dep } x = \infty$.

For arbitrary distinct points $x, y \in \mathbb{R}^n$ we now estimate the expression $h_k(x) - h_k(y)$. We may assume that $\text{dep } x \leq \text{dep } y$. As x and y are distinct, case $\text{dep } y = \infty$ and $\text{dep } x = \infty$ is not possible.

Case 1 Let the Lipschitz constant of the homeomorphism f_i be denoted by λ_i . Let $\lambda = \max\{\lambda_i; 1 \leq i \leq M\}$ and $T = \max\{|G_i|; 1 \leq i \leq M\}$, where $|G_i|$ denotes the number of elements of G_i . Hence the Lipschitz constants of the homeomorphisms $r_1, r_2, \dots, \ell_1, \ell_2, \dots$ do not exceed the number $\Lambda = \lambda^T$. Let $\text{dep } y - \text{dep } x \leq 1$, i.e.

$$\text{dep } x \in \{j, j + 1\}, \quad \text{dep } y = j + 1$$

for some $j \in \mathbb{N} \cup \{-1, 0\}$. By Lemma 2, (iii) and (iv), and because of the construction of h_k ,

$$|h_k(x) - h_k(y)| = |r_{j+1} \circ r_j(x) - r_{j+1} \circ r_j(y)| \leq \Lambda^2 |x - y|.$$

Case 2 Now let $\text{dep } y - \text{dep } x \geq 2$. First let the degrees be nonnegative, i.e.

$$\text{dep } x = j \geq 0 \text{ and } \text{dep } y \geq j + 2$$

for some $j \in \mathbb{N}$. (It may be $\text{dep } y = \infty$ as well.) Then

$$x \in X_{\alpha^j} - \overset{\circ}{X}_{\alpha^{j+1}}, \quad y \in X_{\alpha^{j+2}}.$$

For arbitrary disjoint compact sets $C_1, C_2 \subset \mathbb{R}^n$, set

$$d_{\min}(C_1, C_2) = \min\{|x - y|; x \in C_1, y \in C_2\},$$

$$d_{\max}(C_1, C_2) = \max\{|x - y|; x \in C_1, y \in C_2\}.$$

The sets $X - \overset{\circ}{X}_1$ and X_2 are compact and disjoint; hence the numbers

$$d_X = d_{\min}(X - \overset{\circ}{X}_1, X_2) \quad \text{and} \quad D_X = d_{\max}(X - \overset{\circ}{X}_1, X_2)$$

exist. Since the similarity S_{α^k} maps the triple $(X, X_{\alpha^1}, X_{(\alpha^1, \alpha^2)})$ onto the triple $(X_{\alpha^k}, X_{\alpha^k \alpha^1}, X_{\alpha^k(\alpha^1, \alpha^2)})$, for each $k \in \mathbb{N}$, we have

$$\frac{d_{\max}(X_{\alpha^k} - \overset{\circ}{X}_{\alpha^k \alpha^1}, X_{\alpha^k(\alpha^1, \alpha^2)})}{d_{\min}(X_{\alpha^k} - \overset{\circ}{X}_{\alpha^k \alpha^1}, X_{\alpha^k(\alpha^1, \alpha^2)})} \leq \frac{D_X}{d_X}.$$

Hence $|h_k(x) - h_k(y)| \leq (D_X/d_X) |x - y|$.

Finally, let $\text{dep } x = -1$ and $\text{dep } y \geq 1$, i.e., $x \in \mathbb{R}^n - \overset{\circ}{X}$ and $y \in X_1$. Then $h_k(x) = x$ and

$$\frac{|h_k(x) - h_k(y)|}{|x - y|} \leq \frac{|x - y| + |y - h_k(y)|}{|x - y|} \leq 1 + \frac{\text{diam } X_1}{m},$$

where $m = \inf\{|x - y|; x \in \mathbb{R}^n - \overset{\circ}{X}, y \in X_1\}$ (it is easy to show that $m > 0$). To conclude, set

$$L = \max\left\{\Lambda^2, \frac{D_X}{d_X}, 1 + \frac{\text{diam } X_1}{m}\right\}.$$

Then $|h_k(x) - h_k(y)| \leq L|x - y|$ for an arbitrary $k \in \mathbb{N}$ and $x, y \in \mathbb{R}^n$.

The estimate $|h_k^{-1}(x) - h_k^{-1}(y)| \leq L|x - y|$ can be proved analogously, using Lemma 2(i). \square

4. Main result

Theorem 2. *There exist uncountably many topologically distinct Lipschitz homogeneous wild Cantor sets in \mathbb{R}^3 . In fact, these Cantor sets can all be constructed as simple Antoine Cantor sets with the same number of components of stage n inside each component of stage $n - 1$ and thus with the same Antoine graphs.*

Proof. Use Theorem 1 with $M = 2$, $G_1 = Z_{60}$ and $G_2 = Z_{60}$. Let $\mathbf{T} = (n_1, n_2, \dots)$ be a fixed sequence of 1's and 2's. For each such sequence, construct a Lipschitz homogeneous Antoine Cantor set as in Theorem 1.

For G_1 , let the similarities S_g , $g \in G_1$ be constructed so as to take the outer torus in Figure 1 to the smaller tori in the same figure. Each smaller torus in the chain is obtained from the previous one by rotating the large torus by $2\pi/60$ radians and then by rotating the small tori by $\pi/2$ radians. The homeomorphism f_1 needed in Theorem 1 is constructed in a manner similar to that constructed in the example in [Malešič and Repovš 1999].

For G_2 , let the similarities S_g , $g \in G_2$ be constructed so as to take the outer torus in Figure 2 to the smaller tori in Figure 2. Each smaller torus in the chain is obtained from the previous one by rotating the large torus by $2\pi/60$ radians and then by rotating the small tori by $\pi/4$ radians. The homeomorphism f_2 needed in Theorem 1 is constructed in a manner similar to that constructed in the example in [Malešič and Repovš 1999]. The resulting Cantor set is Lipschitz homogeneously embedded by Theorem 1.

Note that the Antoine graphs associated with any two Cantor set constructed in this way are the same.




Figure 1. Left: A torus with 60 smaller similar tori linked in a simple chain inside, each of which is rotated by $\pi/2$ radians from the previous one. Right: an enlargement of five of the smaller tori.

Let C_1 be the Cantor set constructed as in Theorem 1 for one sequence of 1's and 2's and let C_2 be the Cantor set constructed using a different sequence. We need to show that these two Cantor sets are topologically inequivalently embedded. For this, using the result of Sher mentioned on page 289, it suffices to show that there is no homeomorphism of \mathbb{R}^3 to itself taking the large torus in Figure 1 to the large torus in Figure 2, and taking the chain of smaller tori in Figure 1 to the chain of smaller tori in Figure 2. If there were such a homeomorphism, the link formed by the centerlines of the small tori in Figure 1 would be topologically equivalent to the link formed by the centerlines of the small tori in Figure 2. The next lemma shows that this is not the case.

There are uncountably many sequences of 1's and 2's. The argument above shows that each such sequence leads to a topologically distinct Lipschitz homogeneous wild Cantor set. This completes the proof of the theorem. \square

Lemma 4. *The links formed by the center lines of the smaller tori in Figures 1 and 2 are inequivalent.*

Proof. The link formed by the centerlines of the chain of smaller tori in Figure 2 is seen to be topologically embedded in \mathbb{R}^3 as follows. Starting at a fixed centerline and proceeding around the chain, the centerline can be twisted by a homeomorphism of \mathbb{R}^3 fixed outside of the large torus in such a way that all but one of the centerlines are embedded in the same manner as the link formed by the centerlines of the smaller tori in Figure 1. The remaining centerline in Figure 2 can be obtained from the corresponding centerline in Figure 1 by the following modification. One of the centerlines in Figure 1 is given 30 half twists before linking with the centerlines on either side. A computation of the version of the Jones polynomial introduced in [Kauffman 1988] shows that these links are topologically distinct,




Figure 2. A similar chain, with each small torus rotated $\pi/4$ relative to the previous one.

and thus the Cantor sets are topologically distinct. In fact, all that is needed is to show that the span of the Kauffman polynomial (the highest power appearing in the polynomial minus the lowest power) is different in each case.


For details of a computation of the Kauffman polynomial in this setting, see [Garity \geq 2005]. For completeness, we outline the computation. We refer the reader to [Kauffman 1988] for details on the Kauffman bracket and Kauffman polynomial. Let L represent an oriented link and $|L|$ represent a particular diagram for this link. The writhe of L in the diagram $|L|$ is denoted by $\omega(|L|)$. Let L_+ , L_- , and L_0 represented oriented link diagrams identical to $|L|$ except at one crossing, where L_+ represents this crossing with positive crossing number, L_- with negative crossing number and L_0 with the crossing split and the orientation preserved. We use $X(L)$ to denote the Kauffman polynomial of L .

Open Chains. Let C_n be a simple chain with n links. An easy induction, or the multiplicativity of the Kauffman bracket relative to connected sums, shows that $\langle C_n \rangle = (-A^4 - A^{-4})^{n-1}$, where $\langle \rangle$ is the Kauffman bracket. If we orient the links so the linking numbers alternate signs as in the superscripts below, we have

$$\begin{aligned} X(C_n^{+-\dots+-}) &= X(C_n^{-+\dots-+}) = (-A^4 - A^{-4})^{n-1} && \text{for } n \text{ odd,} \\ X(C_n^{+-\dots-+}) &= (-A)^{-6}(-A^4 - A^{-4})^{n-1} && \text{for } n \text{ even,} \\ X(C_n^{-+\dots+-}) &= (-A)^6(-A^4 - A^{-4})^{n-1} && \text{for } n \text{ even.} \end{aligned}$$

In each case, the span of the polynomial is 8^{n-1} and the maximum and minimum exponent in the polynomial can be read off. The maximum exponent is $4(n - 1)$ if n is odd and $4(n - 1) \pm 6$ if n is even. The minimum exponent is $-4(n - 1)$ if n is odd and $-4(n - 1) \pm 6$ if n is even.

Closed Chains. Now take a closed chain LC_{2n} with $2n$ components and with no twists, as in Figure 1. Orient it so the linking numbers are alternately positive and negative. Consider three consecutive links and modify them as follows:



We now compute using the relation $A^4 X(+) = A^{-4} X(-) + (A^{-2} - A^2) X(\rightleftharpoons)$ and noting that the link labeled L_1 above is $C_{2n}^{+\dots+}$, the link L_{21} is $C_{2n-1}^{-\dots+}$ and the link L_{22} is LC_{2n-2} :

$$\begin{aligned} X(LC_{2n}) &= A^8 X(L_1) + A^4(A^2 - A^{-2})X(L_2) \\ &= A^8 X(C_{2n}^{+\dots+}) + A^4(A^2 - A^{-2})(A^{-8} X(L_{21}) + A^{-4}(A^{-2} - A^2)X(L_{22})) \\ &= A^8 X(C_{2n}^{+\dots+}) \\ &\quad + A^4(A^2 - A^{-2})(A^{-8} X(C_{2n-1}^{-\dots+}) + A^{-4}(A^{-2} - A^2)X(LC_{2n-2})) \\ &= \dots \\ &= (-A^4 - A^{-4})^{2n-2}(-A^6 - A^{-6}) - (A^{-2} - A^2)^2 X(LC_{2n-2}). \end{aligned}$$

Thus

$$X(LC_{2n}) = (-A^4 - A^{-4})^{2n-2}(-A^6 - A^{-6}) - (A^{-2} - A^2)^2 X(LC_{2n-2})$$

This sets up a recursion relation that can be solved for $X(LC_{2n})$. The starting condition is that $X(LC_{2,1}) = (-A^2 - A^{-2})$ has maximum exponent 2 and minimum exponent -2 , and hence span 4.

The maximum exponent of $X(LC_{2n})$ is

$$\max\{4(2n - 2) + 6, 4 + \text{maximum exponent}(X(LC_{2n-2}))\}$$


and the minimum exponent of $X(LC_{2n})$ is

$$\min\{-4(2n - 2) - 6, \text{minimum exponent}((LC_{2n-2})) - 4\}.$$

An easy induction now shows that $\text{span}(X(LC_{2,n})) = 16(n - 1) + 12$ for $n \geq 2$.

Closed chains with twists. Take the case of a linked chain forming a loop with $2n$ components, with k positive half twists, where k is even, in one of the links, LC_{2n}^k .

Orient as in the previous case. By considering the diagram



one can make the following computations.

$$\begin{aligned} X(LC_{2n}^k) &= A^{-8} X(LC_{2n}^{k-2}) + A^{-4}(A^{-2} - A^2)X(C_{2n+1}), \\ X(LC_{2n}^k) &= A^{-8} X(LC_{2n}^{k-2}) + A^{-4}(A^{-2} - A^2)(-A^{-4} - A^4)^{2n}. \end{aligned}$$

The starting point here is when $k = 0$, $LC_{2n}^0 = LC_{2n}$

The maximum exponent is $\max\{\max. \exp.(LC_{2n}^{2k-2}) - 8, 8n - 2\}$.

The minimum exponent is $\min\{\min. \exp.(LC_{2n}^{2k-2}) - 8, -8n - 6\}$.


An easy induction now shows that $\text{span}(X(LC_{2n}^{2k})) = 16n - 4 + 4k$, distinguishing topologically all the chains with $2n$ links and different numbers of even twists. This completes the proof of the lemma. \square

5. Other results and questions

Using techniques similar to those used in the proof of Theorem 1, we can prove the following result. Note that in this case, we assume that G is of the form $Z_p \times Z_q$ for some positive integers p and q .

Theorem 3. *For each i , $1 \leq i \leq 2$, suppose that $f_i: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a Lipschitz homeomorphism and that:*

- (i) $f_i|_{\mathbb{R}^n - \dot{X}} = id$;
- (ii) $f_1(X_{(a,b)}) = X_{(a+1,b)}$ for $(a, b) \in G$;
- (iii) $f_2(X_{(a,b)}) = X_{(a,b+1)}$ for $(a, b) \in G$ and the following diagrams commute:



Then $|(S, T)|$ is Lipschitz homogeneous in \mathbb{R}^n .

The construction suggested by the theorem is similar to the Blankinship construction [1951] for wild Cantor sets in \mathbb{R}^4 .

Question. Can this theorem be used to show that a Lipschitz homogeneous wild Cantor set in \mathbb{R}^4 exists? This would require a more careful Blankinship-type construction, in which the successive stages in the construction were self-similar to the original stage.

Acknowledgment

We thank the referee for helpful suggestions.

References

[Antoine 1920] M. L. Antoine, “Sur la possibilité d’étendre l’homéomorphie de deux figures à leur voisinages”, *C. R. Acad. Sci. Paris* **171** (1920), 661–663. JFM 47.0524.01

[Blankinship 1951] W. A. Blankinship, “Generalization of a construction of Antoine”, *Ann. of Math.* (2) **53** (1951), 276–297. MR 12,730c Zbl 0042.17601

[Eaton 1973] W. T. Eaton, “A generalization of the dog bone space to E^n ”, *Proc. Amer. Math. Soc.* **39** (1973), 379–387. MR 48 #1238 Zbl 0262.57001

- [Garity \geq 2005] D. J. Garity, “Inequivalent Antoine Cantor sets with the same Antoine tree”, in *Proceedings of the Twentieth Annual Workshop in Geometric Topology* (Park City, UT, 2003). To appear.
- [Hutchinson 1981] J. E. Hutchinson, “Fractals and self-similarity”, *Indiana Univ. Math. J.* **30**:5 (1981), 713–747. MR 82h:49026 Zbl 0598.28011
- [Kauffman 1988] L. H. Kauffman, “New invariants in the theory of knots”, *Amer. Math. Monthly* **95**:3 (1988), 195–242. MR 89d:57005 Zbl 0657.57001
- [Malešič and Repovš 1999] J. Malešič and D. Repovš, “On characterization of Lipschitz manifolds”, pp. 265–277 in *New developments in differential geometry* (Budapest, 1996), edited by J. Szenthe, Kluwer, Dordrecht, 1999. MR 99j:57042 Zbl 0939.57023
- [Repovš et al. 1996] D. Repovš, A. B. Skopenkov, and E. V. Ščepin, “ C^1 -homogeneous compacta in \mathbb{R}^n are C^1 -submanifolds of \mathbb{R}^n ”, *Proc. Amer. Math. Soc.* **124**:4 (1996), 1219–1226. MR 1301046 (97f:58008) Zbl 0863.53004
- [Sher 1968] R. B. Sher, “Concerning wild Cantor sets in E^3 ”, *Proc. Amer. Math. Soc.* **19** (1968), 1195–1200. MR 38 #2755 Zbl 0165.57202
- [Wright 1986] D. G. Wright, “Rigid sets in E^n ”, *Pacific J. Math.* **121**:1 (1986), 245–256. MR 87b:57011 Zbl 0697.12018 Zbl 0586.57009

Received April 13, 2004.

DENNIS GARITY
MATHEMATICS DEPARTMENT
OREGON STATE UNIVERSITY
CORVALLIS, OR 97331
UNITED STATES
garity@math.oregonstate.edu
<http://www.math.oregonstate.edu/~garity>

DUŠAN REPOVŠ
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS
UNIVERSITY OF LJUBLJANA
JADRANSKA 19, P.O.BOX 2964
LJUBLJANA
SLOVENIA
dusan.repovs@guest.arnes.si

MATJAZ ŽELJKO
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS
UNIVERSITY OF LJUBLJANA
JADRANSKA 19, P.O.BOX 2964
LJUBLJANA
SLOVENIA
matjaz.zeljko@fmf.uni-lj.si

